



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number: **0 453 242 A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **91303387.4**

(61) Int. Cl.⁵: **A61K 48/00, C12N 15/86**

(22) Date of filing: **16.04.91**

(30) Priority: **16.04.90 US 508731**

(43) Date of publication of application:
23.10.91 Bulletin 91/43

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: **THE GENERAL HOSPITAL
CORPORATION
55 Fruit Street
Boston MA 02114 (US)**

(72) Inventor: **Breakfield, Xandra O.
127 Hovier Street
Newton, Massachusetts 02159 (US)
Inventor: Martuza, Robert L.
31 Fairbanks Road
Lexington, Massachusetts 02173 (US)**

(74) Representative: **Sheard, Andrew Gregory et al
Kilburn & Strobe 30, John Street
London WC1N 2DD (GB)**

(54) **Transfer and expression of gene sequences into central nervous system cells using herpes simplex virus mutants with deletions in genes for viral replication.**

(57) **Methods of gene delivery to cells in the central nervous system are disclosed by introducing and expressing gene sequences mediated by herpes simplex virus 1 (HSV-1) vectors with a mutation in a gene for viral replication.**

EP 0 453 242 A1




FIGURE 1

This invention relates to methods of gene delivery into cells of the central nervous system by introducing and expressing gene sequences using herpes simplex virus 1 (HSV-1) mutants with deletions in gene(s) for viral replication.

The delivery and expression of heterologous or native genes into cells of the nervous system to alter normal cellular biochemical and physiologic processes in a stable and controllable manner is of substantial value in the fields of medical and biological research. This genetic perturbation of the nervous system provides a means for studying the molecular aspects of neuronal function and offering therapeutic approaches to pathologic processes.

Herpes simplex viruses possess several properties that render them attractive candidates as delivery vehicles to bring foreign genes into cells of the peripheral and central nervous systems. (Breakfield, X.O., et al., Mol. Neurobiol. 1:339 (1988); Dobson, A.T., et al., J. Virol. 63:3844-3851 (1989); Ho and Mocarski, Virology 167:279-283 (1988); Ho, D.Y., et al., Proc. Natl. Acad. Sci. USA 86:7596-7600 (1989); Friedmann, T., Science 244:1275-1280; Palella et al., Gene 80:138 (1989).)

These viruses can infect and deliver their DNA into many different cell types, including adult postmitotic neurons; they can enter a state of latency in neurons and possibly other cell types, in which the viral genome exists as an episomal element in the nucleus of the cell; they are able to infect a substantial number of cells *in vivo* due to their ability to reach high titers in culture and to propagate in the nervous system and the relatively long half life of virus particles; large exogenous DNA sequences can be inserted into their genomes; and they possess a wide host range. (Long-necker, R., et al., In: Viral Vectors, Gluzman, Y., et al. (eds.), CSH Lab, pp. 68-72 (1988); Roizmann, B., et al., In: Virology, Fields, B.N., et al. (eds.), Raven Press, New York, pp. 497-526 (1985); Rock, D.L., et al., Nature 302:523-525 (1983).)

Recently, an amplicon-type plasmid-vector system based on herpes simplex virus 1 (HSV-1) has been used to achieve relatively stable expression of the *lacZ* gene in cultured neurons. (Spaete, R.R., et al., Proc. Natl. Acad. Sci. USA 82:694 (1985); Geller, A.I., et al., Science 241:1667 (1988); Geller, A.I., et al., Proc. Natl. Acad. Sci. USA 87:1149 (1990).) In this plasmid vector, the *lacZ* gene was placed under the control of an HSV-1 immediate-early promoter and packaged using a temperature-sensitive HSV-1 helper virus tsK. Coen, D.M., et al., Proc. Natl. Acad. Sci. USA 86:4736 (1989); Leib, D.A., et al., J. Virol. 63:759 (1989); Davison, M.J., et al., J. Gen. Virol. 65:859 (1984).)

Several factors appear to limit the potential uses of herpes viruses for gene transfer to cells in culture and *in vivo*. These include the relatively frequent occurrence of spontaneous revertants of some mutants; possible recombination events between helper virus, viral sequences in plasmids, and latent sequences; deleterious effects on host cell macro-molecular synthesis as a result of viral proteins; alterations in the host cell genome; and possible reactivation of preexisting latent viruses. (Davison, M.J., et al., J. Gen. Virol. 65:859 (1984); Fenwick, M., Compr. Virol. 19:359 (1984); Heilbronn, R., et al., J. Virol. 63:3683 (1989); Kwong, A.D., et al., J. Virol. 63:4834 (1989); Johnson, P.A., et al., 14th International Herpes Workshop abst. (1989).)

In the study of gene expression in neuronal cells in the central nervous system, and the identification of therapies for treating neuronal disease of the central nervous system, a need therefore continues to exist for methods of gene delivery with efficient viral vectors capable of mediating gene transfer into such cells without causing harm to the host animal.

Thus, according to the present invention there is provided a method for introducing and expressing a gene sequence into a central nervous system cell, the method comprising:

introducing into a central nervous system cell a herpes simplex virus 1 (HSV-1) vector comprising a mutation gene for viral replication and a gene sequence operably linked to a promoter sequence so that the gene sequence will be expressed in the cell; and

expressing the gene sequence.

The mutation in the gene for viral replication is preferably in a gene coding for immediate early genes. This may be a gene that encodes infected cell proteins (ICPs) 0, 4, 22, 27 and/or 47.

However, suitably the mutation in the gene for viral replication is in a gene coding for early genes. In such a case, it is preferred that the early genes encodes thymidine kinase or DNA polymerase.

The invention also relates to a pharmaceutical composition comprising the herpes simplex virus 1 mentioned in the above method and a pharmaceutically acceptable carrier.

The invention additionally relates to the use of a herpes simplex virus 1 as described in the above method for use in medicine.

The invention also relates to the use of a herpes simplex virus 1 (HSV-1) vector having a mutation in a gene for viral replication and having a gene sequence operably linked to a promoter sequence, the vector allowing the gene sequence to be expressed in a central nervous system cell so that the expressed gene product complements a neurological deficiency, in the preparation of an agent for treating a neurological deficiency of the central nervous system.

The present invention additionally encompasses the use of a herpes simplex virus 1 (HSV-1) vector with a mutation in a gene for viral replication and having a gene sequence operably linked to a promoter sequence, the vector allowing the expression of the gene sequence in a neuronal cell to modulate neuronal physiology in the preparation of an agent for modulating neuronal physiology in a neuronal cell.

It will be appreciated that the preferred features and characteristics of a method or uses according to the present invention are interchangeable.

The present invention is directed to a method for introducing gene sequences into cells of the central nervous system using HSV-1 mutants as vectors for gene delivery. The HSV-1 virus is mutated so that it has a deletion(s) in a gene(s) for viral replication. The desired gene sequence to be delivered to the cell is inserted into the mutated HSV-1 such that the gene sequence will be expressed in the host cell. By the method according to this invention, a gene sequence is expressed in a cell in the central nervous system by introducing the mutated HSV-1 vector with a mutation in a gene for viral replication and with a gene sequence operably linked to a promoter sequence so that the gene sequence will be expressed in the host cell.

The invention also concerns a method for treating a neurological deficiency of the central nervous system by expressing a gene sequence, which complements the deficiency, in a cell in the central nervous system by introducing a HSV-1 vector with a mutation in a gene for viral replication and with a gene sequence operably linked to a promoter sequence so that the gene sequence will be expressed in the cell and the expressed gene product complements the deficiency.

The invention is further directed to methods for modulating neuronal physiology by delivery of neuropeptide genes, such as those genes that express proteins or poly-peptides to block pain; genes that express growth factors; genes that express proteins or polypeptides to promote regeneration or prolong life-span of a cell; genes that express toxic proteins or polypeptides, for example to kill tumor cells. In the methods for modulating neuronal physiology, the method comprises introducing into a cell a HSV-1 vector with a mutation in a gene for viral replication and with a gene sequence coding for the desired protein or polypeptide operably linked to a promoter sequence so that the gene sequence will be expressed in the cell and upon expression will modulate the neuronal physiology.

Using HSV-1 vectors that have mutations in genes necessary for efficient viral replication, the inventors have discovered methods for introducing and expressing a gene sequence in cells of the central nervous system without toxicity to the host animal.

HSV-1 is a double-stranded DNA virus which is replicated and transcribed in the nucleus of the cell. In virions, HSV-1 vectors are composed of head to tail repeats. (Stow, N.D. *et al.*, Eukaryotic Viral Vectors, Y. Gluzman, Ed. (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1982), pp. 199204; Spaete, R.R. *et al.*, Cell 30:285 (1982).) HSV-1 contains approximately 100 genes and is approximately 150 kilobases in size. Five immediate early ("IE") genes encode infected cell proteins (ICPs) 0, 4, 22, 27, and 47, the major regulatory proteins of the virus. These immediate early proteins then act to regulate the expression of viral proteins of the early (E) and late (L) classes. proteins of the E class are responsible for viral DNA replication. The late ("L") genes are induced after DNA replication and encode the structural components and enzymes required for assembly of virus particles. When the late genes are induced, transcription of the immediate early genes is reduced.

Post-mitotic neurons harbor HSV-1 in the virus' latent state (Stevens, J.G., Curr. Top. Microbiol. Immunol. 70:31-50 (1975).) Once HSV-1 attains latency, it can be retained for the life of the neuron. Latent HSV-1 is capable of expressing genes. Expression of genes encoded by HSV-1 has been detected by *in vitro* hybridization in latently infected neurons. Furthermore, HSV-1 is transported both anterogradely and retrogradely in neurons, and can be passed transsynaptically. These properties are especially advantageous in that it suggests that HSV-1 vectors will be capable of reaching targets of interest some distance away from the injection site.

Two lines of evidence suggest that HSV-1 can infect most, if not all, kinds of neurons in the central nervous system. First, following inoculation of HSV-1 in the periphery, a burst of virus production ascends the neuraxis, initially in the sensory or motor neurons innervating the site of inoculation, then in the spinal cord, brain stem, cerebellum, and cerebral cortex (Koprowski, H., In: Persistent Viruses (Stevens, F.G., ed.), pp. 691-699, Academic Press, NY (1978).) Second, attempts to mimic HSV-1 latency in tissue culture with different preparations of neurons have required high temperature, DNA synthesis inhibitors, and antisera directed against HSV-1 virions to prevent a lytic infection for spreading to all the neurons (Wigdahl, B., *et al.*, Proc. Natl. Acad. Sci. USA 81:6217-6201 (1984).)

In the method according to this invention, HSV-1 vectors with mutations in genes involved in viral replication can be used to effect gene delivery to cells in the central nervous system. As used herein, the term HSV-1 vectors, or mutated HSV-1, is meant to include viral HSV-1 in which the viral sequences have been directly modified so that it may be used for a vector for gene delivery. The term "mutations in genes involved in viral replication" is meant to include gene deletions, gene deficiencies, gene activation, and the like. As stated above, genes

that are involved in viral replication that may be mutated in this invention include the five immediate-early (IE) viral genes encoding regulatory infected cell proteins (ICPs) 0, 4, 22, 27, and 47. These genes are described in detail in Longnecker, R., et al., in: Viral Vectors, Gluzman, Y., et al., (eds.), CSH Lab, pp. 68-72 (1988); Roizmann, B., et al., in: Virology, Fields, B.N., et al., (eds.), Raven Press, New York, pp. 497-526 (1985); Rock, D.L., et al., Nature 302:523-525 (1983).

The early genes that are involved in viral DNA replication that may be mutated and used in this invention include thymidine kinase and DNA polymerase. These early temporal class of expressed HSV-1 genes are described in detail in Longnecker, R., et al., in: Viral Vectors, Gluzman, Y., et al., (eds.), CSH Lab, pp. 68-72 (1988); Roizmann, B., et al., in: Virology, Fields, B.N., et al., (eds.), Raven Press, New York, pp. 497-526 (1985); Rock, D.L., et al., Nature 302:523-525 (1983).

The preferred mutated HSV-1 virus vectors are those that enter latency in neurons and do not reactivate or reactivate very inefficiently.

The present invention concerns a means for in vivo introduction of gene sequences into cells of the central nervous system. The cells of the central nervous system include primarily neurons, but also include neuroglia and other cells. These cells are collectively described herein as "neural or neuronal" cells. Neural cells are described, for example, by Barr, M.L., The Human Nervous System An Anatomic Viewpoint, 3rd. Ed., Harper & Row, NY (1979), which reference is herein incorporated by reference.

The term "gene sequence," as used herein, is intended to refer to a nucleic acid molecule (preferably DNA). Such gene sequences may be derived from a variety of sources including DNA, cDNA, synthetic DNA, RNA, or combinations thereof. Such gene sequences may comprise genomic DNA which may or may not include naturally occurring introns. Moreover, such genomic DNA may be obtained in association with promoter regions or poly A sequences. The gene sequences of the present invention are preferably cDNA. Genomic DNA or cDNA may be obtained in any of several ways. Genomic DNA can be extracted and purified from suitable cells by means well-known in the art. Alternatively, mRNA can be isolated from a cell and used to produce cDNA by reverse transcription, or other means.

As will be appreciated by one of ordinary skill, the nucleotide sequence(s) of the inserted gene sequence or sequences may be of any nucleotide sequence.

The desired gene sequence is operably linked, and under the control of promoter sequence so that the gene sequence will be expressed in the host cell. The promoter sequence is one that is active during viral latency. A viral promoter active during latency can drive stable expression of foreign genes in sensory neurons. (Dobson, A.T., et al., J. Virol. 63:3844-3851 (1989); Ho, D.Y., et al., Proc. Natl. Acad. Sci. USA 86:7596-7600 (1989).) Since herpes virus is known to be maintained in a latent state in the brain, the promoter for the latency-associated transcript (LAT) may be used to mediate stable gene expression. Further, the promoters for the genes for viral replication can be used as promoters to express the gene sequence.

A viral vector, as that term is used herein, is a nucleic acid molecule (preferably of DNA) in which a gene sequence (which is to be transferred) is fused to a subset of viral sequences which are capable of expressing the gene. Also, the viral vector according to this invention will have a mutation in a gene necessary for efficient viral replication. The viral vector should also not have deleterious effects on the host and should infect cells without a helper virus. The viral sequences and the total genome size is selected such that the vector is capable of being encapsulated in a virus particle and thus be capable of binding to, and introducing its gene sequences into a virus-sensitive host cell.

The HSV-1 mutant viral vectors of the present invention can be obtained by transfecting permissive cells with a mixture of plasmid DNA containing HSV-1 gene deletions and infectious and helper HSV-1 DNA. Stow, N.D. et al., Eukaryotic Viral Vectors, Y. Gluzman, Ed. (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1982), pp. 199-204; Spaete, R.R. et al., Cell 30:285 (1982). The obtained mutated HSV-1 viral vectors can be propagated as viruses in appropriate host cells. To grow most mutants, one must transfect a cell with the mutated (and missing) viral gene to replace the deleted viral replication function.

Injection of the vectors to express the desired gene sequence may be done into the brain or into the spinal cord or into nerve endings in the skin, or blood vessels, ventrification surface, etc.

Gene transfer technology has several applications to neuroscience and neurochemistry. The most immediate applications are, perhaps, in elucidating the process of neural peptides and the functional domains of proteins. Cloned cDNA or genomic sequences for neural proteins can be introduced in vivo in order to study cell type-specific differences in processing and cellular fate. By placing the coding sequences under the control of a strong promoter, a substantial amount of the protein can be made, thus avoiding difficulties in characterizing trace amounts. Furthermore, the specific residues involved in protein processing, intracellular sorting, or biological activity can be determined by mutational change in discrete residues of the coding sequence.

Gene transfer technology can also be applied to provide a method to control expression of a protein and to assess its capacity to modulate cellular events in the central nervous system. Certain functions of neural

proteins can be studied *in vivo*, for example, at different times in development or aging in order to monitor changes in receptor density, cell number, fiber growth, electrical activity, and other relevant properties.

Gene transfer provides a means to study the DNA sequences and cellular factors which regulate expression of neural specific genes. One approach to such a study would be to fuse the regulatory elements to be studied to a particular reporter gene and subsequently assaying for the expression of the reporter gene.

The regulation of gene expression in neuronal cells has been found to have a role in maintaining homeostasis and is believed to have a role in mediating information retention in response to external and internal signals (Black, I.B., *et al.*, *Science* 236:1263-1268 (1987).) During development, coordinate regulation of gene expression serves to produce a differentiated phenotype, e.g., as in catecholamine metabolism and myelin biosynthesis. Regulation depends on many factors including chromatin structure, DNA methylation, and transacting factors, which respond to phosphorylation, hormones, and other signals. It is a complex process that allows sets of genes to be expressed together or differentially and may involve a combinatorial code of regulatory sequences.

Issues of cellular fate and interactions in the central nervous system can also be addressed by gene transfer. For example, genes which encode histological markers can be introduced into embryonic cells to determine lineage relationships during development and to elucidate neuronal pathways and to follow cell maturation and fate. In addition, genes encoding growth factors, oncogenic proteins, toxic peptides, or other physiologically important proteins, can be introduced into specific areas of the brain and spinal cord to study their effects on cell division, survival, and differentiation. For some studies, gene transfer or gene expression must be restricted to specific cells in the nervous system.

Gene transfer also possesses substantial potential use in understanding and providing therapy for disease states. There are a number of inherited neurologic diseases in which defective genes are known and have been cloned. In some cases, the function of these cloned genes is known. In humans, genes for defective proteins have been identified for (1) lysosomal storage diseases such as those involving β -hexosaminidase (Komerluk, R.G., *et al.*, *J. Biol. Chem.* 261:8407-8413 (1986); Myerowitz, R., *et al.*, *Proc. Natl. Acad. Sci. USA* 82:7830-7834 (1985)) and glucocerebrosidase (Sorge *et al.*, *Proc. Natl. Acad. Sci. USA* 82:5442-5445 (1985); Tsuji, S., *et al.*, *N. Engl. J. Med.* 316:570-575 (1987)), (2) for deficiencies in hypoxanthine phosphoribosyl transferase activity (the "Lesch-Nyhan" syndrome; Stout *et al.*, *Met. Enzymol.* 151:519-530 (1987)), (3) for amyloid polyneuropathies (prealbumin; Sasaki, H., *et al.*, *Biochem. Biophys. Res. Commun.* 125:636-642 (1984)), (4) for Alzheimer amyloid (Tanzi, R.E., *et al.*, *Science* 235:880-884 (1987); Goldgaber, D., *et al.* *Science* 235:877-880 (1986)); (5) for Duchenne's muscular dystrophy (uncharacterized muscle protein; Monaco, A.P., *et al.*, *Nature* 323:646-650 (1987)); and (6) for retinoblastoma (uncharacterized protein expressed in the retina and other tissues, Lee, W.-H., *et al.*, *Science* 235:1394-1399 (1987); Friend, S.H., *et al.*, *Nature* 323:643-646 (1986).)

Gene transfer techniques can also be used to study the "shiverer" mutation (myelin basic protein, Roach, A., *et al.*, *Cell* 42:149-155 (1987); Molineaux, S.M., *et al.*, *Proc. Natl. Acad. Sci. USA* 83:7542-7546 (1986)) and the "jimpy" mutation (proteolipoprotein, Nave, K.-A., *et al.*, *Proc. Natl. Acad. Sci. USA* 83:9264-9268 (1986); Hudson, L.D., *et al.*, *Proc. Natl. Acad. Sci. USA* 84:1454-1458 (1987).)

The above diseases fall into two classes: deficiency states, usually of enzymes, which are inherited in a recessive manner; and unbalanced states, at least sometimes involving structural or regulatory proteins, which are inherited in a dominant manner.

For deficiency state diseases, gene transfer could be used to bring a normal gene into affected tissues for replacement therapy, as well as to create animal models for the disease using antisense mutations. For unbalanced state diseases, gene transfer could be used to create the disease state in a model system, which could be used in efforts to counteract the effect of the imbalance. Thus, the methods of the present invention permit the treatment of neurological diseases. As used herein, a deficiency state disease is "treated" by partially or wholly remedying the deficiency which causes the deficiency or which makes it more severe. As used herein, an unbalanced state disease is "treated" by partially or wholly remedying the imbalance which causes the disease or which makes it more severe.

The method of this invention may also be used to modulate normal physiologic processes, e.g., delivery of growth factors or other peptides or enzymes to optimize neural regeneration after injury or prolong cell survival in aging or after toxic insults. In addition, it can be used to regulate transmission across certain synapses, for example, to kill neurons in the pathway, or alter neurons within phenotype by up or down regulation of normal neuropeptides or producing neuropeptide analogs as in pain transmission.

In summary, this invention provides a framework for construction and use of replication deficient HSV-1 vectors that can mediate delivery and expression of genes in neurons and other cells in the brain and spinal cord. Virus mutants which cannot replicate in neurons, but can still enter latency, provide the safest vehicle for delivery and appear optimal for targeted delivery to individual neurons either to effect changes in their physiology or to trace their extensions. Mutants which undergo limited replication can be used to alter the physiology

of larger groups of neurons and possibly even through to be transmitted across synapses. Such mutants, however, may be somewhat pathogenic, as cells harbouring productive infections may be killed. Herpes vectors thus offer the potential to alter the physiology of single or groups of neurons to study their function and also provide an avenue of exploration for therapeutic interventions aimed at correcting defects and modulating in neuronal functions.

Having now generally described this invention, the same invention will be described in greater detail by reference to the accompanying drawings and Examples which are included herein for purposes of illustration only and are not intended to be limiting of the invention in the drawings:

Figure 1 shows six photomicrographs illustrating β -galactosidase expression in rat brain cells after injection with herpes simplex virus vectors 7134 and RH105 after 1 day, 3 days, and 14 days;

Figure 2 shows a photomicrograph illustrating β -galactosidase expression in rat brain cells after injection with herpes simplex virus vector GAL4;

Figure 3 is a bar chart showing rat brain areas that contain β -galactosidase expressing cells; and

Figures 4a and 4b show 7134-mediated β -galactosidase gene expression in (a) cerebral cortex, and (b) pyramidal neurons in cerebral cortex, respectively.

EXAMPLE A

Three mutant viruses were studied to explore the potential of these replication deficient mutants as vectors to deliver genes in CNS cells *in vivo*.

The first vector, 7134, provided by Drs. Priscilla Schaffer and Weizhong Cai (Dana Farber Cancer Center) possesses lacZ substitutions in both copies of HSV-1 ICPO gene. The construction of this vector is described in detail in Cai, W., et al., *J. Virol.* 63:4579 (1989), incorporated herein by reference. Expression of lacZ in this mutant is regulated by the ICPO immediate-early promoter. Compared to wild-type HSV-1, ICPO mutants replicate poorly in a variety of cells in culture and *in vivo*. These mutants establish latency in sensory neurons of trigeminal ganglia as frequently as wild-type HSV-1, when used to inoculate mouse corneas. Once they have established latency, however, they reactivate very inefficiently. (Leib, D.A., et al., *J. Virol.* 63:759-768 (1989); Sacks, W.R., et al., *J. Virol.* 61:829-839 (1987).)

The second vector, GAL4, provided by Dr. Neal DeLuca, possesses lacZ substitutions in both copies of the ICP4 gene. This vector is described in detail in Shepard, A.A., et al., *J. Virol.* 63:3714 (1989), incorporated herein by reference. Expression of lacZ is controlled by the early promoter for the ICP6 gene. ICP4 mutants do not replicate in culture or when inoculated onto mouse corneas. As a consequence, ICP4 mutants are unable to establish latency efficiently, and cannot reactivate from the latency state. (DeLuca, N.A., et al., *J. Virol.* 62:732-743 (1988); Leib, D.A., et al., *J. Virol.* 63:759-768 (1989).)

The third vector, RH105, provided by Drs. Edward Mocarski and Dora Ho (Stanford) contains a lacZ substitution in the thymidine kinase gene. This vector is described in detail in Ho, D.Y., et al., *Virology* 167:279 (1988), incorporated herein by reference. Expression of this gene is regulated by the ICP4 immediate-early promoter. Thymidine kinase (TK) gene is described in palella et al., *Gene* 80:138 (1989); Coen, D.M., et al., *Proc. Natl. Acad. Sci. USA* 86:4736 (1989); Ho, D.Y., et al., *Virology* 167:279 (1988). TK mutants can replicate in actively growing cells in culture (presumably by utilizing the endogenous cellular thymidine kinase), but replicate poorly following inoculation peripherally or in nervous system cells into rodents. (Leist, T.P., et al., *J. Virol.* 63:4976 (1989); Coen, D.M., et al., *Proc. Natl. Acad. Sci. USA* 86:4736 (1989).) These mutants can enter latency in sensory neurons of peripheral ganglia, but do not reactivate from the latent state. (Leist, T.P., et al., *J. Virol.* 63:4976 (1989); Coen, D.M., et al., *Proc. Natl. Acad. Sci. USA* 86:4736 (1989).)

Previous animal studies with similar mutant viruses have shown that they are relatively nonpathogenic. (Ho, D.Y., et al., *Virology* 167:279 (1988); Coen, D.M., et al., *Proc. Natl. Acad. Sci. USA* 86:4736 (1989); Leib, D.A., et al., *J. Virol.* 63:759 (1989).) However, these experiments were based on inoculation of epidermal tissues (corneas or footpads) and histological assessment of infections in the sensory ganglia of the peripheral nervous system.

These three vectors, 7134, GAL4, and RH105, were used to express the *Escherichia coli lacZ* gene in neuronal cells of rat central nervous system. These vectors possessed lacZ deletional substitutions in the genes encoding the immediate-early viral proteins, ICP0 and ICP4, and the early protein, thymidine kinase, and consequently were compromised or defective in their ability to replicate. In all cases lacZ was placed under immediate-early or early viral promoters active in the early phase of infections. Expression of β -galactosidase was observed in cortical neurons following stereotactic inoculation of these mutant viruses into adult rat brains. All mutant vectors exhibited markedly reduced pathogenesis in animals as compared to wild type HSV-1 KOS strain. Different patterns of expression of the β -galactosidase gene were observed with the different vectors used. Injection of the ICP0 mutant in frontal cortex and caudate produced β -galactosidase expression in a sub-

stantial number of cells around the inoculation site and at some distance from it for up to 14 days. The ICP0 vector appeared to have been transported retrogradely to the contralateral cingulate cortex. The ICP4 and thymidine kinase mutants showed only a few cells expressing β -galactosidase cells in brain areas immediately adjacent to the injection tract for a few days after injections. These herpes virus-derived vectors provide an opportunity for the in situ delivery and expression of specific genes in neurons in the central nervous system (CNS) with little adverse effect on animals.

Example 1

In order to evaluate the potential use of vectors 7134, GAL4, and RH105 for direct gene delivery into CNS cells, the potential pathogenic effects on animals following direct intracerebral inoculations was first assessed. Using 200,000 plaque forming units (PFUs) of 7134, GAL4, RH105, or the wild-type KOS strain of HSV-1 in 2 μ l volumes, these vectors were injected into the right frontal area of adult rat brains. As a control, equal volumes of medium alone were injected. Rat survival and behavior were then assessed daily for two weeks. Table I shows the survival results for these animals. While all animals survived for three days without apparent abnormalities, seven of eight rats injected with KOS and one of seven rats injected with 7134 exhibited profound behavioral abnormalities. The rats injected with KOS showed either lethargy or marked hyperactivity after 3 days, had frequent seizures within 4-5 days, and died by 6 days. The animal that did not survive the 7134 injection was lethargic and showed a marked decrease in feeding and drinking activities starting approximately 8 days after injection and its death occurred rapidly thereafter. One animal that received an intracerebral injection of 7134 became lethargic and perished after 8 days. At autopsy, foci of β -galactosidase positive neurons were found dispersed throughout the cerebral cortex, particularly in the temporal lobes. This pattern appeared similar to the one seen in the rats that had received intracerebral injections of the wild-type HSV-1 KOS. Since this particular rat had been housed mistakenly in a cage that contained rats which had been injected with wild-type virus, one possible explanation for the widespread distribution of lacZ positive cells in this animal is that it also became infected with wild-type HSV-1 KOS, presumably through some of the blood and serum fluid found at the craniotomy site. This might have created a helper effect allowing for the widespread replication of 7134 in this animal.). Control animals and those injected with GAL4 or RH105 viruses showed no behavioral abnormalities or death over a two-week period. These results indicate that these HSV mutant vectors, GAL4, RH105, and 7134, are relatively nonpathogenic and that behavioral abnormalities do not routinely result from their use in rats. A repeated study with 7134 did not show the anomaly previously exhibited.

Example 2

After inoculations as described in Example 1, rats were sacrificed one, three and fourteen days after virus inoculations in order to assess the extent of β -galactosidase expression. Brains were processed histochemically and assessed for the presence of β -galactosidase activity. In the presence of the substrate X-gal at alkaline pH, this enzyme yields an intense blue stain throughout the cytoplasm of infected cells. (Turner, D.L., et al., Nature 328:131 (1987); Price, J., et al., Proc. Natl. Acad. Sci. USA 84:156 (1987); Sanes, J.R., et al., EMBO J. 5:3133 (1986).)

With the aid of a stereotactic apparatus, approximately 200,000 PFUs of each mutant virus were injected into the right frontal area of rat brains (stereotactic coordinates: AP +2.7/LAT -0.3/Depth -0.4). Two μ l volumes were injected with a Hamilton syringe using stereotactic coordinates. Rats were then monitored daily for abnormalities in feeding and drinking behavior and for level of activity. One, three, and fourteen days after viral injections, animals were anesthetized and sacrificed by intracardiac perfusion with 200 ml of 3% paraformaldehyde in phosphate buffered saline. Anesthesia was obtained by intraperitoneal injection of 0.5 ml-1 ml of a solution that contained 16.2% nembutal, 10% ethanol, 39.6% polyethylene glycol, 166 mM magnesium sulfate and 4.25% chloral hydrate. Brains were then processed over the following 3 days by serially placing them for approximately 24 hours in the same solution which contained increasing concentrations of sucrose (15%, 20%, and 30%). After freezing on dry ice and storage at 70°C, brains were sectioned on a cryostat in 40 μ m sections. After washing with phosphate buffered saline, sections were reacted overnight in a solution containing 0.1% X-Gal, 2 mM magnesium chloride, 35 mM potassium ferrocyanide, 35 mM potassium ferrocyanide, 0.1% sodium deoxycholate and 0.1% NP4.0. After washing in phosphate buffered saline, sections were mounted on glass slides and examined by light microscopy.

Numerous cells expressing β -galactosidase were observed one, three and 14 days after stereotactic injection of 7134 into the right front cortex. While most cells expressing β -galactosidase were localized within a 175 μ m² and around the injection site at day 1, positive cells were found in a 600 μ m² area by day 3. By day 14 there was a diminution in the number of cells expressing β -galactosidase with sparse foci located (100 μ m²)

around the needle track.

In contrast to the findings for 7134, one and three days after RH105 injection, sparse foci of β -galactosidase expressing cells were present only along the needle track (covering an area of $10 \mu\text{m}^2$), and by 14 days no positive cells could be detected (Figure 1).

Similarly, after GAL4 injection, only an occasional cell expressing β -galactosidase were observed (Figure 2). Procedures were the same as those given for Figure 1, except that, after the β -galactosidase histochemical reaction, the brain sections were dehydrated in 70% and 80% ethanol and then counterstained with neutral red. Sections were then rinsed in 80% ethanol and water before examining them by light microscopy.

These findings indicate that two patterns of gene delivery are characteristic of these vectors. The first, represented by 7134, is that of a replication compromised vector capable of delivering a gene to a relatively large number of cells in the CNS with limited spread away from the site of a delivery. The second, represented by GAL4 and RH105, is that of a replication defective vector that delivers a gene to only a few cells at the injection site without spread to other sites.

The differences in the patterns of gene expression in the CNS among the vectors could also reflect the inactivation of different viral genes, or genetic differences amongst the viral strains used. Vectors 7134 and GAL4 were obtained from an HSV-1 KOS virus strain that had been propagated in one laboratory, whereas RH105 was derived from an HSV-1 KOS strain propagated in a different laboratory. The wild-type HSV-1 KOS strain used as a control in all experiments was also obtained from the same laboratory that vectors 7134 and GAL4 were obtained.

The differences in the patterns of gene expression in the CNS among the vectors may also reflect variations in promoter strength. This latter possibility is not likely, however, since all three promoters, ICP0 (7134), ICP6 (GAL4), and ICP4 (RH105), are expressed with immediate-early kinetics during productive infection.

The pattern of 7134-mediated β -galactosidase expression is best explained by an initial productive infection in numerous cells near the injection site followed by secondary spread of viral particles to adjacent brain areas. Eventual limit to the spread of this virus could be mediated by decreasing yields of virus from infected cells and entry into latency, and/or by immunosuppression. By contrast, the pattern of RH105 or GAL4-mediated β -galactosidase expression indicates that only a small number of cells near the injection site are infected initially, and that no productive infection occurs, although the virus may enter latency.

As controls, brains from rats that had been injected in right frontal areas with the wild-type virus and with medium alone were examined. A monoclonal antibody ID-4 against the major capsid protein of the virus was used to detect infected neurons in rats five days after injection with wild-type HSV-1 KOS. At this time these animals exhibited profound behavioural abnormalities.

Brain sections from HSV-1 KOS injected rats were rinsed in Tris-buffered-saline (TBS) solution (pH 7.4) and incubated for one hour with horse serum (diluted according to Vectastain ABC kit specifications). The sections were then reacted for one hour with medium harvested from the ID-4 monoclonal antibody producing hybridoma line. After thorough rinsing in TBS, the sections were incubated for one hour with biotinylated horse anti-mouse IgG (Vectastain). After additional rinsing, the sections were placed for 30 minutes in Vectastain ABC solution, containing avidin and then in a solution containing 0.01% hydrogen peroxide and 0.05% diaminobenzidine tetrahydrochloride in 0.1 M Tris HCP, pH 7.2 for 2-5 minutes. Sections were then rinsed in TBS and mounted on glass slides.

The immunohistochemical reaction detected numerous areas where viral antigen was present at a distance from the injection site (results not shown). These included bilateral cortical regions in the frontal, cingulate, temporal and parietal lobes. This suggested that the limited number of β -galactosidase expressing cells seen following injection of the mutants is a property of the mutants and is not due to technical artifacts. When medium alone was injected, no blue staining was visible even at higher magnification (results not shown). This control was necessary due to reports that the β -galactosidase histochemical reaction will at times detect endogenous lysosomal β -galactosidase activity. Ho, D.Y., et al., *Virology* 167:279 (1988).

Figure 3 represents rat brain areas that contain β -galactosidase expressing cells. Areas along the needle track were calculated by measuring the furthest linear distance between cells expressing β -galactosidase in two perpendicular directions and then multiplying the two values. Several sections (range: 3-11) were measured for each inoculation from at least two different rats (range: 2-4). Error bars indicate S.D. from mean). Bar graphs represent the averages of these measurements. Measurements were taken with the aid of an ocular micrometer on a light microscope.

To determine which type of cells expressed β -galactosidase, following inoculation of 7134 into the right caudate, sections were then examined for β -galactosidase expression after counterstaining with neutral red. Both small and large types of caudate neurons, as identified by morphology, expressed β -galactosidase three days later.

Stereotactic injections of 7134 were performed in the right caudate nucleus (stereotactic coordinates: AP

+1.7; LAT -0.3; Depth -0.4). Animals were sacrificed three days later. Brains were then processed, stained for β -galactosidase activity and counterstained with neutral red as described in the preceding figures.

In some rats expression of the lacZ gene was evident in a relatively large area of the cingulate cortex (Figure 4a), probably due to retrograde transport of the virus from the caudate. Examination of the cingulate cortex at higher magnification revealed discrete labeling of pyramidal neurons, identified by characteristic morphology (Figure 4b). Reaction product was apparent throughout dendrites and axons. Most of these neurons did not show evidence of degeneration, such as swelling or beading of processes. It was also evident β -galactosidase gene expression was not as apparent in non-neuronal cell types.

In these studies, lacZ gene transfer and expression into CNS neurons was achieved using replication deficient HSV-1 mutants. The majority of cells expressing lacZ following inoculation of the 7134 replication-compromised mutant possessed the morphological characteristics of neurons. In addition, many of the neurons expressing lacZ, (following caudate inoculation) were found some distance away in the cingulate cortex, raising the possibility that the virus possesses a particular tropism for this cortical region. This finding also supports the occurrence of retrograde transport of the mutant virus following from nerve endings in the caudate to cell bodies in the cortex. Transneuronal transport can deliver genes to specific neurons via inoculation into areas containing their axonal or dendritic projections. Recent findings using wild type HSV-1 indicate that transneuronal and transsynaptic transfer can be followed using ^3H -thymidine labeling and immunocytochemical staining of viral antigens. (Ugolini, G., et al., *Science* 243:89 (1989); Norgren, Jr., R.B., et al., *Brain Res.* 479:374-378 (1989); Margolis, T.P., et al., *J. Virol.* 63:4756-4761 (1989); Kuypers, G.J.M., et al., *TINS* 13:71-75 (1990).) The viral vectors with the lacZ substitutions thus should be used for transsynaptic transport. This provides a useful tool to neuroanatomically trace specific neuronal projections and synaptic pathways with less toxicity to animals.

TABLE 1. SURVIVAL OF RATS AFTER INTERCEREBRAL INJECTION OF VIRUS

| <u>Days After Injection</u> | | | |
|-----------------------------|--------------------|-------|------------------|
| <u>Mutant</u> | 1 | 3 | 14 |
| Control ^a | 12/12 ^b | 8/8 | 4/4 |
| 7134 | 26/26 | 8/8 | 6/7 ^c |
| 7134 | 12/12 | 12/12 | 12/12 |
| GAL4 | 15/15 | 11/11 | 3/3 |
| RH105 | 12/12 | 8/8 | 4/4 |
| KOS | 19/19 | 15/15 | 1/8 ^d |

^aEqual volume of medium was injected.

^bExpressed as animals surviving/animals inoculated on day 0.

^cDeath of one animal occurred 8 days after injection. This animal exhibited lethargy and inactivity prior to death.

^dAnimal deaths occurred 6-7 days after inoculation and were preceded by frequent seizures 5 days following inoculation.

Claims

1. A method for introducing and expressing a gene sequence into a central nervous system cell, the method comprising:
 5 introducing into a central nervous system cell a herpes simplex virus 1 (HSV-1) vector comprising a mutation gene for viral replication and a gene sequence operably linked to a promoter sequence so that the gene sequence will be expressed in the cell; and
 expressing the gene sequence.
- 10 2. A method according to claim 1 wherein the mutation in the gene for viral replication is in a gene coding for immediate early genes.
3. A method according to claim 2 wherein the gene coding for immediate early genes is a gene that encodes infected cell proteins (ICPs) 0, 4, 22, 27 and/or 47.
- 15 4. A method according to claim 1 wherein the mutation in the gene for viral replication is in a gene coding for early genes.
5. A method according to claim 4 wherein the gene coding for early genes encodes thymidine kinase or DNA
 20 polymerase.
6. The use of a herpes simplex virus 1 (HSV-1) vector having a mutation in a gene for viral replication and having a gene sequence operably linked to a promoter sequence, the vector allowing the gene sequence
 25 to be expressed in a central nervous system cell so that the expressed gene product complements a neurological deficiency, in the preparation of an agent for treating a neurological deficiency of the central nervous system.
7. The use of a herpes simplex virus (HSV-1) vector having a mutation in a gene for viral replication and having a gene sequence operably linked to a promoter sequence, the vector allowing expression of the gene sequence
 30 in a neuronal cell to modulate neuronal physiology in the preparation of an agent for modulating neuronal physiology in a neuronal cell.
8. The use according to claim 6 or 7 wherein the mutation in a gene for viral replication is in a gene coding for an immediate early gene, such as an immediate early gene which encodes infected cell proteins (ICPs)
 35 0, 4, 22, 27, or 47.
9. The use according to claim 6 or 7 wherein the mutation is a deletion in a gene coding for early genes.
10. The use according to claim 9 wherein the gene coding for early genes encodes thymidine kinase or DNA
 40 polymerase.




FIGURE 1

EP 0 453 242 A1




FIGURE 2



FIGURE 3




FIGURE 4a

EP 0 453 242 A1



FIGURE 4b



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 91 30 3387

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|--|---|--|---|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (Int. CL.5) |
| X, D | GENE. vol. 80, no. 1, August 1989, AMSTERDAM NL pages 137 - 144; PALELLA, T.D. et al.: "Expression of human HPRT mRNA in brains of mice infected with a recombinant herpes simplex virus-1 vector" * the whole document * | 1, 4-7 | A61K48/00 C12N15/86 |
| X | WO-A-9002551 (BIOSOURCE GENETICS CORPORATION) * the whole document * | 1, 4-7 | |
| P, X | WO-A-9009441 (THE GENERAL HOSPITAL CORPORATION) * the whole document * | 1, 4-7 | |
| P, X | THE NEW BIOLOGIST vol. 2, no. 8, August 1990, pages 739 - 746; CHIOCCA, E.A. et al.: "Transfer and expression of the lacZ gene in rat brain neurons mediated by Herpes Simplex Virus mutants" * the whole document * | 1-9 | |
| | | | TECHNICAL FIELDS SEARCHED (Int. CL.5) |
| | | | A61K C12N |
| The present search report has been drawn up for all claims | | | |
| Place of search THE HAGUE | | Date of completion of the search 15 JULY 1991 | Examiner CHAMONNET F.J. |
| <p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application I : document cited for other reasons & : member of the same patent family, corresponding document</p> | | | |

EP 0 FORM 1503 (01.91) (P0401)

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)